

NKR46 pico 8: involvement of network in treatment of anorexia

Review information

Authors

Danish Health Authority¹

[Empty affiliation]

Citation example: DHA. NKR46 pico 8: involvement of network in treatment of anorexia. Cochrane Database of Systematic Reviews [Year], Issue [Issue].

Characteristics of studies

Characteristics of included studies

Dare 2001

Methods	Participants	Interventions	Outcomes	Notes

Risk of bias table

Bias	Authors' judgement	Support for judgement
Sequence generation	Low risk	From Fisher 2010
Allocation concealment	Low risk	From Fisher 2010
Blinding of participants and personnel	High risk	From Fisher 2010
Incomplete outcome data	Unclear risk	From Fisher 2010
Selective reporting	High risk	From Fisher 2010
other bias	Unclear risk	From Fisher 2010

Russell 1987 group 2 and 3

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias table

Bias	Authors' judgement	Support for judgement
Sequence generation	Unclear risk	From Fisher 2010
Allocation concealment	Low risk	From Fisher 2010
Blinding of participants and personnel	Unclear risk	From Fisher 2010
Incomplete outcome data	High risk	From Fisher 2010
Selective reporting	High risk	From Fisher 2010
other bias	High risk	From Fisher 2010

*Footnotes***Characteristics of excluded studies*****Ball 2004***

Reason for exclusion

Crisp 1991

Reason for exclusion

Eisler 2000	Reason for exclusion
Espina 2000	Reason for exclusion
Geist 2000	Reason for exclusion
Hall 1987	Reason for exclusion
Ie Grange 1992	Reason for exclusion
Lock 2005	Reason for exclusion
Rausch 2006	Reason for exclusion
Robin 1999	Reason for exclusion
Russell 1987	Reason for exclusion

Russell 1987 group 1

Reason for exclusion

Russell 1987 group 2

Reason for exclusion

Whitney unpublished

Reason for exclusion

Footnotes

Characteristics of studies awaiting classification

Footnotes

Characteristics of ongoing studies

Footnotes

References to studies

Included studies

Dare 2001

Published and unpublished data

Russell 1987 group 2 and 3

Excluded studies

Ball 2004

Published data only (unpublished sought but not used)

[Empty]

Crisp 1991

Published data only (unpublished sought but not used)

[Empty]

Eisler 2000

Published and unpublished data

[Empty]

Espina 2000

Published data only (unpublished sought but not used)

[Empty]

Geist 2000

Published data only (unpublished sought but not used)

[Empty]

Hall 1987

Published data only (unpublished sought but not used)

[Empty]

Ie Grange 1992

Published and unpublished data

[Empty]

Lock 2005

Published data only (unpublished sought but not used)

[Empty]

Rausch 2006

Published data only (unpublished sought but not used)

[Empty]

Robin 1999

Published and unpublished data

[Empty]

Russell 1987

Published and unpublished data

[Empty]

Russell 1987 group 1

Published and unpublished data

[Empty]

Russell 1987 group 2

[Empty]

Whitney unpublished

Unpublished data only

[Empty]

Data and analyses

1 Active involvement vs no active involvement of network in treatment

Outcome or Subgroup	Studies	Participants	Statistical Method	Effect Estimate
1.1 ED behavoir, longest FU	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
1.2 ED behavoir, ECT	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
1.3 Weight, follow-up (5 years)	1	32	Mean Difference (IV, Random, 95% CI)	1.00 [-7.65, 9.65]
1.3.1 %ABW	1	32	Mean Difference (IV, Random, 95% CI)	1.00 [-7.65, 9.65]

NKR46 pico 8: Involvement of network in treatment of anorexia

14-Jan-2021

1.4 Weight post intervention	1	32	Mean Difference (IV, Random, 95% CI)	3.00 [-5.39, 11.39]
1.4.1 %ABW	1	32	Mean Difference (IV, Random, 95% CI)	3.00 [-5.39, 11.39]
1.5 Psychological symptoms, follow-up	1	32	Mean Difference (IV, Random, 95% CI)	-1.20 [-3.04, 0.64]
1.5.1 family based therapy	1	32	Mean Difference (IV, Random, 95% CI)	-1.20 [-3.04, 0.64]
1.6 Psychological symptoms, post intervention	1	35	Mean Difference (IV, Random, 95% CI)	-0.30 [-2.09, 1.49]
1.6.1 Family based therapy	1	35	Mean Difference (IV, Random, 95% CI)	-0.30 [-2.09, 1.49]
1.7 Maintenance of remission (follow-up)	1	33	Risk Ratio (M-H, Random, 95% CI)	1.46 [0.80, 2.67]
1.7.2 5 years	1	33	Risk Ratio (M-H, Random, 95% CI)	1.46 [0.80, 2.67]
1.8 Dropout	2	80	Risk Ratio (M-H, Random, 95% CI)	1.67 [0.62, 4.52]
1.8.2 during therapy	2	80	Risk Ratio (M-H, Random, 95% CI)	1.67 [0.62, 4.52]
1.9 Quality of life, longest FU	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable
1.10 Wellbeing of care-givers, longest FU	0	0	Mean Difference (IV, Fixed, 95% CI)	Not estimable